

- 01** Wound inside the mouth will be healed 2-3 weeks later. Beware of wound opening during this period. It is safe to eat soft food, not brushing teeth, and it is recommended not to mouth inflate like a balloon when gargling.
- 02** Take antibiotics for 5 days in order to prevent inflammation. Also, it is important to keep mouth clean after consuming a meal.
- 03** Bleeding may occur within 24-48 hours after surgery. Square jaw reduction, two jaw surgery, one jaw surgery and protruding mouth surgery apply to this matter. Patient will be hospitalized with compression bandage on the day of surgery. If bleeding is suspected on the next day of surgery, patient cannot be discharged. Patient can be discharged when bleeding is not suspected after 24 hours, however, complete rest as well as observation from the guardian is necessary for 1-2 days after discharge. If face swells up with worsening pain, please contact hospital immediately.
- 04** There could be change in sensation of facial skin after surgery. Usually, it returns to normal after several months, however, rarely it could last continuously. Patient can experience sensation of teeth pulling, stiff mouth and various other feelings in the beginning after surgery, however, mostly it disappears without problem.
- 05** It needs 2-3 months for union of fractured cheekbone after cheekbone reduction surgery. During this period, be careful of impact or pressure to the cheekbone area. Also, refrain from opening mouth wide, chewing hard food, and clenching mouth severely.
- 06** Both sides of the face cannot be perfectly symmetrical. From the head, two sides are slightly different and complete symmetry cannot be achieved even after surgery.
- 07** In case of any questions or emergency, please contact (+82-10-7156-6546).